

**Microplastic pollution in the coastal water
of the Peter the Great Gulf:
content and distribution.
The first stage of survey**

NOWPAP POMRAC

Vladivostok, Russia
2017

*Pollution Monitoring Regional Activity Center of UNEP Action Plan for the Protection,
Management and Development of the Marine and Coastal Environment
of the Northwest Pacific Region (NOWPAP POMRAC)*

*Региональный Центр по мониторингу загрязнения окружающей среды Плана действий
ЮНЕП по охране, управлению и развитию морской и прибрежной среды в регионе
северо-западной Пацифики (NOWPAP POMRAC)*

*Pacific Geographical Institute, Far Eastern Branch of the Russian Academy of Science
Федеральное Государственное Бюджетное Учреждение Науки
Тихоокеанский Институт Географии Дальневосточного отделения РАН*

Microplastic pollution in the coastal water of the Peter the Great Gulf: content and distribution. The first stage of survey / Authors: N.V. Kozlovskii, A.N. Kachur. Editor: S.Yu. Moninets.– Vladivostok: PGI FEB RAS, 2018. – 44 p.

Загрязнение прибрежных вод залива Петра Великого микропластиком: его состав и распределение. Первый этап исследований / Авторы: Н. В. Козловский, А.Н. Качур. Отв. Редактор: С.Ю. Монинец. – Владивосток: ТИГ ДВО РАН, 2018. – 44 с.

For bibliographic purposes, this document should be cited as follows:

Kozlovskii N.V., Kachur A.N. (2018) Report on microplastic pollution in the coastal water of the Peter the Great Gulf: content and distribution. The first stage of survey. POMRAC Technical Report N 13. – Vladivostok: PGI FEB RAS, 2018. – 44 p.

Table of contents

1. <u>Introduction</u>	5
1.1. <u>Background</u>	5
1.2 <u>Objectives of the activity</u>	5
2. <u>Selection of sampling sites</u>	6
3. <u>Methods applied in this study</u>	8
3.1. <u>Sampling methods</u>	8
3.1.1. <u>Hand net sampling</u>	8
3.1.2. <u>Neuston net sampling</u>	9
3.2. <u>Sample treatment procedure</u>	10
3.2.1. <u>Drying of the sample</u>	11
3.2.2. <u>Removal of natural organic matter</u>	11
3.2.3. <u>Density separation and filtering</u>	12
3.3. <u>Type/size description of microplastics</u>	13
3.4. <u>Weight measurement</u>	15
3.5. <u>Polymer type identification</u>	15
4. <u>Distribution of plastic particles</u>	16
4.1. <u>Khasan seashore (site 1)</u>	17
4.2. <u>Cape Nazimov (site 2)</u>	18
4.3. <u>Minonosok Inlet (site 3)</u>	19
4.4. <u>Srednyaya Bight (site 4)</u>	20
4.5. <u>Slavyanka Bay (site 5)</u>	21
4.6. <u>Perevoznaya Bight (site 6)</u>	21
4.7. <u>Peschany Peninsula (site 7)</u>	22
4.8. <u>Chaika beach (site 8)</u>	22
4.9. <u>Steklyannaya Bight (site 9)</u>	25

<u>4.10. Lazurnaya Bay (site 10)</u>	<u>25</u>
<u>4.11. Strelok Bay (site 11)</u>	<u>26</u>
<u>4.12. Nakhodka Bay (site 12)</u>	<u>26</u>
<u>5. Discussion</u>	<u>27</u>
<u>5.1. Hotspots</u>	<u>27</u>
<u>5.2. Morphological composition</u>	<u>27</u>
<u>5.3. Size differentiation</u>	<u>29</u>
<u>5.4. Polymer composition</u>	<u>29</u>
<u>5.5. Comparison of hand net sampling results and results of neuston trawling</u>	<u>30</u>
<u>5.6. Comparison of microplastic concentrations in the study area and worldwide</u>	<u>32</u>
<u>5.7. Suggested seasonal factor and land-based sources</u>	<u>33</u>
<u>6. Conclusion</u>	<u>38</u>
<u>7. Acknowledgements</u>	<u>40</u>
<u>References</u>	<u>41</u>

1. Introduction

1. 1. Background

Plastic contamination of the marine and coastal areas has become one of the most important aspects of marine litter issue around the world. The highest concern is raised about the microplastic, i.e. small-sized fragments or unfragmented items with size <5 mm. The marine microplastic is proved to cause harm to the aquatic organisms (P. Davison et al., 2011; F. Murray et al., 2011) and seabirds (M. D. Robards et al., 1995), through both direct harm of ingestion (plastic particles, mistaken by food, may cause indigestion in adult individuals and underdevelopment in young organisms and reproductive disruption (Sussarellu et al., 2016)) and contamination with associated chemicals, such as POPs or plasticizers, involving these substances into the trophic chains (Teuten et al., 2009). Therefore, understanding of the situation with microplastic pollution in the NOWPAP region, which is one of the most densely populated areas in the world, is necessary. The NOWPAP regional countries, including Japan, People's Republic of China and Republic of Korea already carry out their national marine microplastic studies, while in the Russian Federation this activity is new. Understanding microplastic contamination levels in the coastal areas is very important because they sustain commercial near-shore fisheries and aquaculture. This study provides new and valuable information about the potential sources, distribution pattern and composition of microplastic pollution in the Northwest Pacific region.

1. 2. Objective of the activity

This project was performed as a part of Regional Action Plan on marine Litter (RAP MALI) activity of NOWPAP POMRAC for the biennium of 2016-2017 and focused on a survey of microplastic pollution in the southern coastal area of the Russian Far East.

The overall objective of the study was to estimate concentrations of microplastic particles in the coastal seawater of the study area, describe their type/size characteristics, distribution, and possible sources. In the two selected

areas (mainly in the coastal area near the Tumen River mouth and a beach near Vladivostok) we conducted seasonal surveys to understand the spatio-temporal variations in the distribution of contaminants within a year. We also compared the number and weight of microplastics in the tidal zone, which is a primary target in this study regarding finding possible land-based sources of pollution, with measurements conducted in the nearby water areas located 100-300 meters off the coast.

Prior to starting this activity, preliminary studies were carried out to prove that microplastic pollution is a threat to the coastal ecosystems of the Russian NOWPAP area (Blinovskaya, 2016; Yakimenko and Blinovskaya, 2016; unpublished data by PGI of August 2015). Considering that this project was an initial attempt to assess the microplastic contamination in the study area and had a limited budget, the research has to be continued to have a better understanding of microplastics pollution in the Russian Far East.

The results of the study, besides highlighting the pollution hazard, are directly related to the activity of NOWPAP POMRAC on setting Ecological Quality Objectives (objective 5: “Marine litter does not adversely affect coastal and marine environments”). According to the structure of the EcoQOs, further studies of microplastic contamination should include impacts of litter on marine life by monitoring of trends in the amount and composition of litter ingested by marine animals. Patterns of coastal microplastic pollution, especially in registered hot spots, could be useful for developing spatial criteria assessing contamination of aquatic life with plastic litter.

2. Selection of sampling sites

The Peter the Great Gulf is the largest gulf (9,000 km²) in the Sea of Japan and consists of three larger bays, i.e. Amur Bay, Ussuri Bay, and Possiet Bay, and smaller bays, such as Nakhodka, Strelak, and etc. The Amur Bay coastal area is one of the most populated areas in Primorsky Krai. Most part of Vladivostok city located on the Muravyov-Amurskiy peninsula, the home of over 600 thousand

people is located along the Amur Bay. The coasts of the Ussuri Bay, in the eastern part of the Muravyov-Amurskiy peninsula, and the Nakhodka Bay are also relatively densely populated areas compared to other coasts of Primorsky Krai.

The sampling sites (Fig. 1) are mainly chosen based on the criteria of population density and remoteness (as a baseline), though factors of seasonal recreation and river discharge were also considered. Urban coastal sites are located along the eastern part of the Amur Bay, while along the western part, smaller populated areas (including those used for summer recreation and area near the Razdolnaya/Suifen River mouth) are common.

In the Possiet Bay, water area of marine biosphere reserve, and the coast of Khasansky District, are located near the Tumen River mouth. In the Ussuri Bay, Nakhodka Bay, and Strelok Bay, summer recreation areas, urban areas, and sites near river mouths were sampled.



Fig. 1. Sampling sites: 1) – Khasan Seashore; 2) Cape Nazimov; 3) Minonosok Inlet; 4) Srednyaya Bight; 5) Slavyanka Bay; 6) Perevoznaya; 7) Peschany Peninsula; 8) Chaika Beach; 9) Steklyannaya; 10) Lazurnaya Bay; 11) Strelok Bay; 12) Nakhodka Bay

3. Methods

In this study, the most appropriate and affordable methodology of sampling was applied.

The sampling and analysis methods used are similar to those applied by other NOWPAP members, especially Japan (Tokyo University of Marine Science and Technology, Kyushu University) and Republic of Korea (Korean Institute of Ocean Studies and Technologies).

Most samples in this study were collected in the tidal zone (0-5 m off the coastline). Several seawater samples were collected at the distance of 100-300 m from the shore, mainly in the areas of regular survey, to register possible differences in microplastic concentrations depending on the distance from the coastline. Two different methods were applied to the two different settings.

3.1.1. Hand net sampling

In this study, we used a plankton net for the sampling in the tidal zone (see table XXXX showing the basic parameters of this net). It is transformed into a hand-net by attaching a retractable rod (the adjusted length is approx. 1-3 meters) to its mouth ring (fig 2 a). The samples are collected along the selected sampling site by horizontal filtering with the net half-submerged into the water (fig. 2 c) To calculate the volume of filtered water, we applied a mechanical flowmeter (Hydro-bios, model 438-110, Germany) attached to the net mouth (fig. 2 b). The volume of filtered water is calculated based on the number of the impeller rotations and the submerged net mouth area. The samples are transferred into plastic bottles of 1000 ml. The volume of filtered water in the tidal zone depends to a considerable extent on the amount of suspended sand and algae at the sampling location. After each filtering, the net walls are rinsed at least twice from the outside into the sample container to collect remaining plastic particles. After each sampling procedure, the net is rinsed in fresh water.



Fig. 2. Hand net sampling. Left to right – a) the net with a retractable rod, b) the flowmeter, c) sampling

3.1.2. Neuston net sampling

A neuston net was used to collect water samples at the distance 100-300 m from the shoreline. Table 1 shows the basic parameters of this net.

The manta net/neuston net is considered (Hidalgo-Ruz et al., 2012) as a highly efficient tool for horizontal water sampling in sea. It is very useful for filtering of tens or hundreds cubic meters of water (depending on the mouth area) for a relatively short time (10-20 minutes). The construction of manta net provides its high buoyancy without a need to control the depth of its mouth frame, while the depth of neuston net should be controlled during the tow. We had a neuston net for the research. During the first sampling, the net was directly towed by a motor boat (fig. 3). Later, we attached it (with additional frame providing adjustable fixed depth) to a catamaran dragged by a motor boat. In this case, suddenly reduced towing speed did not result in the submerging of the net frame below sea surface (fig. 4). The average speed of the trawling was 1.5 knots, and the time was 15-20 minutes or more. The water volume was also calculated based on the number of revolutions of flowmeter attached under the boat/catamaran.



Fig. 3. Neuston net sampling.
Net directly towed by boat



Fig. 4. Neuston net sampling. Net with adjustable frame is attached to a catamaran

As in case with hand net samples, the filtered neuston trawl was poured into plastic bottles, including water used for rinsing the net walls.

After collection, the samples were brought to the laboratory for further analysis.

Net type	Mouth dimensions, m	Overall length, m	Opening area, m ²	Mesh size, mm
Plankton net	Diameter: 0.2	0.4	0.03	0.1
Neuston net	Width: 0.5 Depth: 0.2	1	0.1	0.1

Table1. Basic parameters of plankton/neuston net we used

3.2. Sample treatment procedure

The sample treatment was mainly based on the protocols adopted from the Korean Institute of Ocean Science and Technology (KIOST), especially in dealing with natural organic matter contained in the samples. It generally coincides with the NOAA sample treatment protocols (Laboratory Methods..., 2015). The following stages are included in the process:

3.2.1. Drying of the sample.

In the laboratory, the collected samples are filtered with a mesh size 0.1 mm and the filtered samples are transferred into the beakers (fig. 5). Spatula and minimal amount of distilled water are applied in the process, after that the sample is covered with holed foil to avoid airborne contamination and dried at temperature of 25 C° (ambient temperature) to 60 C° (using desiccator).



Fig.5. Left- a sample in a beaker before drying, right- a dried sample being weighed

3.2.2. Removal of natural organic matter.

After drying, 20 ml of 0.05M Fe(II)SO₄ water solution is applied to the sample, then 30% H₂O₂ is added (20mg or more, if necessary) by small amounts to remove extra organic matter. In this reaction, Fe(II) is used as a catalyst to accelerate the process of decomposition. After most of organic matter is decomposed by the reaction, 6 grams of NaCl is added per each 20 ml of the sample. After adding NaCl, the sample is carefully mixed with a glass rod to make the solution more homogeneous (fig. 6).



Fig.6. Left - adding hydrogen peroxide; right - NaCl

3.2.3. Density separation and filtering

The next step is transferring of the solution in the density separator funnel, including careful rinsing of the beaker that contained the sample (fig. 7). After that, the funnel is covered with foil to avoid contamination and settled for a day to deposit heavy solid fraction. Next, the lower fraction is drained from the funnel to check for possible microplastic particles remaining and then it is discarded. The supernatant and particles remaining on the funnel walls are then carefully washed with distilled water into the filtering system. The filters applied are 47mm diameter polycarbonate filters with a pore size of 5 μm . After filtering, the particles remaining on the walls of filtering system are carefully transferred to the filter paper using forceps and distilled water. Then filters are dried at room temperature in petri dishes for further analysis.



Fig.7. After density separation in a funnel, samples are transferred to polycarbonate filters.

3.3. Type/size description of microplastics

Sorting and type-size identification of the plastic particles is carried out using a stereomicroscope (Discovery V12). By size classification, the following major groups are selected: 0.1-1 mm (smaller microplastics), 1-5 mm (larger microplastics), and 5-25 mm (mesoplastics). In this research, the size of the smallest microplastic particles is 0.1 mm corresponding to the sampling mesh size. The size of the plastic particles is determined using the camera application installed on the microscope. Main types of the obtained plastic are identified as follows: fibers (fig. 10 a), fragments (fig. 8), films (fig. 9), and foam (fig. 10 b). We also registered paint fragments and microbeads (fig.11), but as of now, their number tends to be low in the sites of survey.





Fig.8. Fragments





Fig.9. Films



Fig. 10. a) Fibers; b) Foam (EPS) on 0.3 mm steel mesh




Fig.11. a) microbead (diameter 0.1 mm); b) paint fragment (diameter 0.6 mm)

3.4. Weight measurement

During the type/size identification the plastic-like objects are retrieved from each sample and transferred to preliminarily weighed envelopes made of tracing paper for determining their weight with analytical scale. Before weighing, the envelopes with and without plastics are dried under 80degrees C for 4 hours to remove extra water contained by the tracing paper.

3.5. Polymer type identification

Identification of polymers (fig.12) is conducted with application of attenuated total reflection spectroscopy (ATR) using Diamond ATR device set on the Fourier transform infrared spectrometer (Thermo Nicolet 6700). The spectrum of each identified object is determined based on a series of scans ($\Sigma = 32$).




Fig.12. FTIR spectrometer and IR-spectra of the most frequent polymer types in the marine environment (Polyethylene(PE), Polypropylene (PP), and Polystyrene(PS)). Reference spectra of each polymer type are depicted below actual spectra of marine microplastics.

4. Distribution of plastic particles

Fig. 13 shows generalized distribution of microplastics among the selected sampling sites of the coastal area of the Peter the Great Gulf. The data is compiled for summer and fall seasons in 2016 and 2017 across 12 sampling sites. After processing the initial sampling data, the focus of our further survey was on the sites showing higher concentrations of microplastic particles. These sites are the coastal area adjoining to the Tumen River mouth and the eastern part of the Amur Bay.




Fig. 13. Average distribution of floating micro- and mesoplastic particles in the tidal zone of the Peter the Great Gulf in July-September 2016-2017 (particles in m³).

Description of each selected site, including information on the geographical location, composition of floating microplastics, possible sources of contamination, seasonal peculiarities, etc. is provided below.

4.1. Khasan seashore (Site 1)

This area is located far from inhabited localities and is difficult to assess due to its remoteness. No considerable economic activities, except seasonal bird hunting, are registered there. Tourists and beach goers are basically not allowed in that area without a permission because it is neighboring to the state border. We chose this site for the survey because it is very close to the Tumen River mouth (approximately 5 km from the estuary). The Tumen River is an important transboundary water body shared by the three countries, including China, North Korea, and Russia. Even though the river is not used for shipping since 1940, its basin is densely populated (by approximately 2 million people in China and North Korea). The river may serve as a source of microplastic pollution in the selected area. The sampling site is located on a wide and long sandy beach (approximately 10-15 km in length) stretching from the river estuary to the Sivuchya Bay, which belongs to the Far Eastern Marine Biosphere Reserve. We collected three samples at this location: in September 2016, in July 2017, and in October 2017. Each sample displayed high concentrations of micro- and meso plastic fragments in the tidal zone. As of now, these figures considerably exceed floating micro-litter concentrations from all other sampling sites selected in the study.



Fig.14. Beached EPS on the Khasan shore (28 September 2016)

Sampling date, dd/mm/yy	Sample volume, m ³	Basic fragment types, p./m ³		Polymer types	Number, p./ m ³	Weight, mg/m ³
27/09/2016	2.5	Fragment	12	PE, PP, PS	54	15.8
		Film	21	PE, PP		
		Fiber	11	Nylon, Polyester, PP		
		Foam	10	PS		
28/07/2017	3.6	Fragment	23	PE, PP, PS	56	57.6
		Film	19	PE, PP		
		Fiber	6	Nylon, Polyester, PP		
		Foam	8	PS		
11/10/2017	2	Fragment	10	PE, PP	31	6.9
		Film	14	PE, PP		
		Fiber	5	Nylon, Polyester		
		Foam	2	PS		
Average number/weight±SD					47±14	26.7±27.1

Table 2. Khasan seashore, description of floating micro- and mesoplastics in the tidal zone

Besides the fact that the concentrations of floating plastic particles are higher than elsewhere in this study, number of EPS particles was not less than 8p./m³ in all three samples (14-25%), which is specific to this area. The beach is covered with dense patches of foam spherules or larger EPS fragments (fig. 14), and we calculated that beached EPS micro-and mesoplastic could reach 95-99% of all other plastic fragments at this site.

4.2. Cape Nazimov (Site 2)

This site is located in the apex of the Posyet Bay, northeast to the previously described sampling site. This area borders with few inhabited localities and with the port of Possiet (approx. 4,500 people). Cape Nazimov is a popular recreation place in summer, though it is uninhabited. We suggested that the water area could be polluted with microplastics considering its relative vicinity from the highly littered site. Nevertheless, microplastic concentrations there tend to be low (0.5-2 p./m³).

We collected two samples from the area, in September 2016 and in June 2017. During the recreational season, the concentrations turned to be even lower than in late September. It is difficult to accurately assess which fragment types prevail there due to low total number of floating microplastic particles in both samples (approx. 5 m³ contained 15 microplastic particles)

Sampling date, dd/mm/yy	Sample volume, m ³	Basic fragment types, p./m ³		Polymer types	Number, p./ m ³	Weight, mg/m ³
29/09/2016	2.4	Fragment	1	PE	3	0.1
		Film	2	PE, PP		
		Fiber	-	-		
		Foam	-	-		
28/07/ 2017	2.7	Fragment	-	-	3	0.05
		Film	2	PE, PP		
		Fiber	1	Polyester		
		Foam	-			
Average number/weight					3	0.075

Table 3. Cape Nazimov, description of floating micro- and mesoplastics in the tidal zone

4.3. Minonosok Inlet (site 3)

The Minonosok Inlet is a small water area in the Posyet Bay, which is a part of the western section of the Far Eastern Marine Biosphere Reserve (FEMBR). This area is aquaculture farms for the cultivation of such commercial species as scallop and oyster. These farms provide for the restoration of natural scallop population in the reserve.

Sampling date, dd/mm/yy	Sample volume, m ³	Basic fragment types, p./m ³		Polymer types	Number, p./ m ³	Weight, mg/m ³
04/06/2016	0.9	Fragment	21	PE, PP	36	11.9
		Film	9	PE, PP, PTFE		
		Fiber	5	Nylon, PE		
		Foam	1	PS		

Table 4. Minonosok Inlet, description of floating micro- and mesoplastics in the tidal zone

Sampling date, dd/mm/yy	Sample volume, m ³	Basic fragment types, p./m ³		Polymer types	Total number, p./ m ³	Weight, mg/m ³
04/06/2016	15	Fragment	0.4	PE	0.7	n/a
		Film	0.3	PE, PP		
		Fiber	-	-		
		Foam	-	-		

Table 4.1. Minonosok Inlet, description of floating micro- and mesoplastics collected by neuston trawl

We had a chance to join the cruise of R/V Nasonov in late May-early June 2016 and collected samples in Minonosok Inlet (table 4 and 4.1.) and in Srednyaya Bight (table 5 and 5.1.), which both belong to the FEMBR. We cannot describe seasonality impact in the inlet because we collected only one sample there, but evidently weathered PE buoys on the shore considerably litter the water area with microplastics.

4.4. Srednyaya Bight (site 4)

Sampling date, dd/mm/yy	Sample volume, m ³	Basic fragment types, p./m ³		Polymer types	Number, p./ m ³	Weight, mg/m ³
05/06/2016	1.1	Fragment	5	PE	11	0.5
		Film	4	PE, PP		
		Fiber	2	Nylon		
		Foam	-	-		

Table 5. Srednyaya Bight, description of floating micro- and mesoplastics in the tidal zone

Sampling date, dd/mm/yy	Sample volume, m ³	Basic fragment types, p./m ³		Polymer types	Total number, p./ m ³	Weight, mg/m ³
05/06/2016	15	Fragment	0.13	PE, PP	0.4	n/a
		Film	0.27	PE, PP		
		Fiber	-	-		
		Foam	-	-		

Table 5.1. Srednyaya Bight, description of floating micro- and mesoplastics collected by neuston trawl



Fig. 15. Left - Srednyaya Bight, right- Peschany Peninsula

4.5. Slavyanka Bay (site 5)

The area we chose for sampling is a tourist beach exposed to an intense flow of tourists during summertime. Local population in adjoining areas is relatively small (around 12,000 people), therefore the site could have been characteristic for understanding a seasonal input of microplastics (especially of primary type). We collected samples in late September 2016 and in July 2017 (table 6).

Sampling date, dd/mm/yy	Sample volume, m ³	Basic fragment types, p./m ³	Polymer types	Number, p./ m ³	Weight, mg/m ³	
28/09/2016	2.2	Fragment	-	-	0.5	<0.01
		Film	-	-		
		Fiber	0.5	Polyester		
		Foam	-	-		
28/07/ 2017	2	Fragment	0.5	PE	5.5	0.2
		Film	1	PE, PP		
		Fiber	4	Polyester		
		Foam	-			
Average number/weight				3	0.1	

Table 6. Slavyanka Bay, description of floating micro- and mesoplastics in the tidal zone

4.6. Perevoznaya Bight (site 6)

This water area is located near a settlement with a small population of less than 1,000 people.

Sampling date, d/m/y	Sample volume, m ³	Basic fragment types, p./m ³		Polymer types	Total number, p./ m ³	Weight, mg/m ³
28/09/2016	2	Fragment	0.5	PE	1.5	0.05
		Film	0.5	PE		
		Fiber	0.5	Polyester		
		Foam	-	-		

Table 7. Perevoznaya Bight, description of floating micro- and mesoplastic in the tidal zone

4.7. Peschany Peninsula (site 7)

This area is sparsely populated, though local people use the adjoining water for fishing. Also, it is located opposite to the Vladivostok city. The table below (table 8) shows the results of sampling we carried out in September 2016 and in July 2017.

Sampling date, d/m/y	Sample volume, m ³	Basic fragment types, p./m ³		Polymer types	Total number, p./ m ³	Weight, mg/m ³
28/09/2016	3	Fragment	-	-	3.5	0.5
		Film	3	PE, PP		
		Fiber	-	-		
		Foam	0.5	PS		
26/07/ 2017	4.7	Fragment	1.5	PE	6.7	2.03
		Film	5	PE, PP		
		Fiber	0.2	Polyester		
		Foam	-	-		
Average number/weight					5.1	1.3

Table 8. Peschany Peninsula, description of floating micro- and mesoplastics in the tidal zone

4.8. Chaika beach (site 8)

This is a narrow pebble-covered beach (approx. 10 meters in width) located on the western coast of the Amur Bay. This site is exposed to urban impacts as it is located in close proximity to Vladivostok city. Also, it is used for summer tourism and recreational fishing. The location is near to the Pacific Geographical Institute and easy to assess. We carried out more detailed seasonal monitoring in the tidal

Date, d/m/y	Vol., m ³	Fr. types, p./m ³		Polym. types	N-r., p./ m ³	W-t, mg/m ³
21/07/2016	2	Fragment	1	PE	11	2.1
		Film	7.5	PE, PP		
		Fiber	2	PP, Polyester		
		Foam	0.5	PS		
16/02/2017	3	Fragment	-	-	0.3	<0.001
		Film	-	-		
		Fiber	0.3	Polyester		
		Foam	-	-		
16/05/2017	3	Fragment	7	PE	15.6	2.25
		Film	2.3	PE, PP		
		Fiber	6.3	Polyester, PE, PP		
		Foam	-	-		
18/06/2017	2.5	Fragment	3.2	PE, PP	14.4	2.08
		Film	3.2	PE, PP		
		Fiber	8	Polyester, PP		
		Foam	-	-		
16/07/2017	3	Fragment	2	PE	12	1.9
		Film	3	PE, PP		
		Fiber	7	Polyester		
		Foam	-	-		
08/08/2017	3.1	Fragment	2	PE	17.5	2.1
		Film	7.5	PE, PP		
		Fiber	8	Polyester, nylon		
		Foam	-	-		
16/09/2017	3	Fragment	0.5	PE	7.5	0.2
		Film	1	PE		
		Fiber	6	Polyester, nylon		
		Foam	-	-		
18/10/2017	2.8	Fragment	-	-	5	0.1
		Film	1	PE		
		Fiber	4	Polyester		
		Foam	-	-		
15/11/2017	3	Fragment	-	-	5.3	0.03
		Film	0.3	PE		
		Fiber	5	Polyester, nylon		
		Foam	-	-		
Average number/weight					9.8	1.2

Table 9. Chaika Beach, description of floating micro- and mesoplastics in the tidal zone

zone, and compared summer data for the tidal zone to neuston trawling data (usually in July-August). The table 9 shows the results of tidal water sampling in 2016-2017.

Besides seasonal sampling in the tidal zone, we collected several samples by trawling. In general, ratio of plastics collected by neuston net trawl to those collected by the hand net was approximately 10 times lower.

In August 2015 we collected preliminary samples by towing of a plankton net in that area to find any plastic fragments in the seawater to substantiate the necessity of this research. Despite the fact that various tools were applied (neuston net with rectangular mouth and opening area of 0.1 m² against plankton net with circular mouth and opening area of 0.03 m²), the collected data seems to be very important showing high concentrations of fibers in that area (table 10)

Sampling date, d/m/y	Volume, m ³	Basic fragment		Polymer types	N-r., p./ m ³	W-t, mg/m ³
		types, p./m ³				
12/08/2015	≈120*	Fragment	0.01	PE, PS	1.7	n/a
		Film	0.05	PE, PP		
		Fiber	1.6	n/a		
		Foam	-	-		
		Microbead	0.01	PE		
21/07/ 2016	47	Fragment	0.25	PE	1.1	0.08
		Film	0.14	PE, PP		
		Fiber	0.71	Polyester		
		Foam	-	-		
		Microbead	-	-		
08/08/2017	65	Fragment	0.18	PP, PE, PS	0.6	0.05
		Film	0.12	PE, PP		
		Fiber	0.22	Polyester, PE		
		Foam	0.08	PS		
		Microbead	0.01	PE		
Average number/weight					1.13	0.06

Table 10: Chaika beach, description of floating micro- and mesoplastics collected by neuston trawl

*We did not use flowmeter in August 2015. The volume of filtered water was calculated by the formula: $V=Qt$, where Q is water discharge and t is time.

Q was calculated as follows: $Q=vS$, where v is the speed of the boat, and S is the net opening area.

4.9. Steklyannaya Bight (site 9)

This is a bight located in the Ussuri Bay which is also exposed to impacts of Vladivostok city. This site is also used for summer recreation. We collected tidal zone samples and neuston samples (tables 11 and 12) to compare concentrations in two largest bays of the gulf (i.e. the Amur Bay and the Ussuri Bay).

Sampling date, dd/mm/yy	volume, m ³	Basic fragment types, p./m ³		Polymer types	N-r, p./ m ³	W-t, mg/m ³
20/06/2017	3.2	Fragment	2.7	PE, PP, PS	11	0.3
		Film	2.7	PE, PTFE		
		Fiber	5.6	Polyester, PE, PP		
		Foam	-	-		
08/08/2017	3	Fragment	1.7	PE, PP	10.7	0.25
		Film	2	PE, PP		
		Fiber	7	Polyester		
		Foam	-	-		
Average number/weight					10.8	0.27

Table 11. Steklyannaya Bight, description of floating micro- and mesoplastics in the tidal zone

Sampling date, d/m/y	volume, m ³	Fragment types, number, p./m ³		Polymer types	N-r, p./ m ³	W-t, mg/m ³
08/08/2017	62	Fragment	0.01	PE	0.13	0.005
		Film	-	-		
		Fiber	0.12	Polyester, PE		
		Foam	-	-		

Table 12. Steklyannaya Bight, description of floating microplastics collected by neuston trawl

4.10. Lazurnaya Bay (site 10)

Lazurnaya Bay is a part of the Ussuri Bay. This is a wide and long sandy beach with strongly expressed seasonal human load, being the most popular recreational site in the vicinity of Vladivostok city. We sampled this site in October 2017 and in July 2017. The results of sampling are shown below (table 13).

Sampling date, d/m/y	Sample volume, m ³	Fragment types, number, p./m ³		Polymer types	N-r, p./ m ³	W-t, mg/m ³
20/10/2016	3	Fragment	1	PE, PP, PS	9	0.1
		Film	1	PE, PTFE		
		Fiber	7	Polyester, PE, PP		
		Foam	-	-		
09/07/2017	3.1	Fragment	2	PE, PP, PS	15.3	0.8
		Film	3.3	PE, PP		
		Fiber	9.7	Polyester, PP		
		Foam	0.3	PS		
Average number/weight					10.8	0.27

Table 13. Lazurnaya Bay, description of floating micro/mesoplastics collected in the tidal zone

4.11. Strelok Bay (site 11)

It is a bay in the eastern part of the gulf. The population in the adjoining area is much lower than along eastern coast of the Amur Bay and western coast of the Ussuri Bay, though this area is used for summer recreation. Only one sample was collected in autumn. The results of sampling are shown below (table 14).

Sampling date, d/m/y	volume, m ³	Fragment types, number, p./m ³		Polymer types	N-r, p./ m ³	W-t, mg/m ³
04/09/2017	2	Fragment	2.5	PE, PP	5	0.4
		Film	0.5	PE		
		Fiber	2	Polyester		
		Foam	-	-		

Table 14. Strelok Bay, description of floating micro- and mesoplastics in the tidal zone

4.12. Nakhodka Bay (site 12)

The Nakhodka bay washes the city of Nakhodka and adjoining port area. We collected only one sample in the autumn, so there is a need for more detailed seasonal study in this area. Table 15 shows the results of sampling.

Sampling date, d/m/y	volume, m ³	Fragment types, number, p./m ³		Polymer types	N-r, p./ m ³	W-t, mg/m ³
04/09/2017	3	Fragment	1	PE	4	0.1
		Film	-	-		
		Fiber	3	Polyester, PP		
		Foam	-	-		

Table 15. Nakhodka Bay, description of floating micro- and mesoplastics in the tidal zone

5. Discussion

5.1. Hotspots

The results of our survey show two areas with elevated concentrations of microplastics in the tidal zone of the Peter the Great Gulf - coastal area near Tumen River estuary and the adjoining waters in the southern part of the Posyet Bay and coastal area around the Muravyov-Amursky Peninsula (eastern part of the Amur Bay and western part of the Ussuri Bay) close to the city of Vladivostok. Microplastic contamination in the tidal zone of Minonosok Inlet (south-eastern part of the Posyet Bay) is also high, reaching 36 particles per m³. Near Vladivostok urban area, the eastern coast of the Amur Bay and the western coast of the Ussuri Bay are exposed to microplastic contamination to approximately similar extent, reaching 11-16 particles per m³ and 10-15 particles per m³, respectively.

The lowest concentrations of microplastics in the tidal zone are registered in the northern part of the Posyet Bay and in the Perevoznaya Bay (3 particles per m³ and 1.5 particles per m³). The highest concentrations of microplastics in the offshore seawater were measured along the coastline of Khasan district (Khasan Seashore), reaching 56 plastic particles of various shape and size per m³.

5.2. Morphological composition

The composition of microplastics in the tidal zone differs for each location. In number, all morphological types obtained from the Khasan area (near Tumen River estuary) prevail or comparable to other sampling locations. Number of fragments

there is comparable to Minonosok Inlet, but considerably overcomes all other sites (6-20 times). Number of films overcome all sampling locations (2.5–20 times). Number of fibers in the tidal zone of Khasan is comparable to concentrations registered near the Vladivostok agglomeration, but also higher than elsewhere in this survey. Foamed styrene concentrations are relatively low in the gulf, except Khasan area (up to 10 times higher than in other sampling locations).

Sampling site	Fragment, Number/m ³		Film, Number/m ³		Fiber, Number/m ³		Foam, Number/m ³	
	min-max	(%)	min-max	(%)	min-max	(%)	min-max	(%)
1) Khasan shore	12-23	(22-41%)	14-21	(39-34%)	6-11	(11-20%)	8-10	(14-19%)
2) Nazimov Cape	0-1	(0-33%)	9	(67%)	0-1	(0-33%)	0	(0%)
3) Minonosok Inl.	21	(58%)	4	(25%)	5	(14%)	1	(3%)
4) Srednyaya B.	5	(46%)	0-1	(36%)	2	(18%)	0	(0%)
5) Slavyanka Bay	0-0.5	(0-9%)	0.5	(0-18%)	0.5-4	(73-100%)	0	(0%)
6) Perevoznaya B.	0.5	(33.3%)	3-5	(33.3%)	0.5	(33.3%)	0	(0%)
7) Peschany Pen.	0-1.5	(0-22%)	2.3-7.5	(75-86%)	0-0.2	(0-3%)	0-0.5	(3-22%)
8) Chaika Beach	1-7	(22-44%)	2-2.7	(14-22%)	2-8	(39-56%)	0-0.5	(0-3%)
9) Steklyannaya	1.7-2.7	(16-19%)	1-3.3	(19-25%)	5.6-7	(51-53%)	0	(0%)
10) Lazurnaya	1-2	(11-13%)	0.5	(11-22%)	7-9.7	(63-78%)	0-0.3	(0-2%)
11) Strelok Bay	2.5	(44.5%)	0	(11%)	2	(44.5%)	0	(0%)
12) Nakhodka B.	1	(25%)	2	(0%)	3	(75%)	0	(0%)

Table 16. Minimum and maximum concentrations of microplastic particles by morphological type

Table 16 shows minimum and maximum concentrations of microplastics by

morphological type, though for many sites this distribution may be disputable due to low total number of microplastic particles collected or only one sampling made. The table does not include winter samples because only one site was selected for winter sampling (Chaika Beach).

5.3. Size differentiation

For most of the sampling sites the fraction of 0.1 – 1 mm was more abundant, ranging from approximately 55 to 30-35 percent, followed by the 1-5 mm fraction (45-25 percent). Mesoplastics was found mainly in the most contaminated areas, for example near the Tumen River estuary (fig. 17).




Fig. 17. Size range of floating microplastic particles in the tidal zone of Khasan seashore, July 2017 (0.1-1 mm – smaller microplastics, 1-5 – larger microplastics, 5-25 – mesoplastics)

5.4. Polymer composition

Regarding polymer type composition, in the most reliable (considering the total number of microplastics) sampling sites prevailing is polyethylene (50-70%) followed by polypropylene (16-36%) and polystyrene (8-25%). Fig. 18 shows distribution of polymers in key sampling locations. The data for polyester and nylon should be interpreted with caution because they constituted less than 10% of total fibers and no FTIR-microscopy was applied for precise identification. Among the microfiber samples that we managed to identify, about 50% were made of polyester,

the other 15-20% were nylon and the rest were PP and PE, and two fibers were PS.




Fig. 18. Distribution of basic polymer types in key sampling locations

5.5. Comparison of hand net sampling results and results of neuston trawling

Unlike hand net samples, we collected neuston samples only in summer. That is why we can compare summer neuston trawls (used in offshore sampling) and summer hand net samples (used in tidal zone sampling) (table 17). We conducted neuston net trawling in four locations, including Minonosok Inlet, Srednyaya Bight, eastern part of the Amur Bay (near Chaika Beach), and western part of Ussuriisky Bay (Steklyannaya Bight). The table below shows differences between concentrations in the tidal zone and 100-300 meters off the coast. The results indicate that microplastic concentrations drastically decrease from the intertidal zone to the open water by an order of magnitude. In the absence of more frequent measurements, it is impossible to conclude if the decrease is linear or there is an abrupt change along the transect from the tidal zone to the offshore.

Sampling location	Sampling method	Sampling date, d/m/y	Fragment types, number, p./m ³		N-r, p./ m ³	W-t, mg/ m ³
1) Minonosok Inlet	Hand net sampling	04/06/2016	Fragment	21	36	11.9
			Film	9		
			Fiber	5		
			Foam	1		
	Neuston trawling	04/06/2016	Fragment	0.4	0.7	n/a
			Film	0.3		
			Fiber	-		
			Foam	-		
2) Srednyaya Bight	Hand net sampling	05/06/2016	Fragment	5	11	0.5
			Film	4		
			Fiber	2		
			Foam	-		
	Neuston trawling	05/06/2016	Fragment	0.13	0.4	n/a
			Film	0.27		
			Fiber	-		
			Foam	-		
3) Chaika beach (eastern part of the Amur Bay)	Hand net sampling	16/07/2017	Fragment	2	12	1.9
			Film	3		
			Fiber	7		
			Foam	-		
	Neuston trawling	08/08/2017	Fragment	0.18	0.6	0.05
			Film	0.12		
			Fiber	0.22		
			Foam	0.08		
Microbead			0.01			
4) Steklyannaya Bight (western part of the Ussuri Bay)	Hand net sampling	08/08/2017	Fragment	1.7	10.7	0.25
			Film	2		
			Fiber	7		
			Foam	-		
	Neuston trawling	08/08/2017	Fragment	0.01	0.13	0.005
			Film	-		
			Fiber	0.12		
			Foam	-		

Table 17. Comparison of hand net /neuston net samples




Fig. 19. An example of varying concentrations of 3 types of plastic fragments from tidal zone to 300 meters seawards (Srednyaya Bight, June 2016)

5.6. Comparison of microplastic concentrations in the study area and worldwide

Location	Mesh (μm)	MP (par. /m ³)	References
SE Black Sea	200	600–1200	Aytan et al., 2016
China, Yangtze Estuary	32	500–10200	Zhao et al., 2014
China, Yangtze Estuary	333	0.03–0.455	Zhao et al., 2014
SE Korea, coast	50	592–1299	Kang et al., 2015
SE Korea, coast	330	4.22–44.28	Kang et al., 2015
Sweden, coast	80	150–2400	Norén, 2008
NE Pacific (off British Columbia CA)	62.5–250	8–9180	Desforges et al., 2014
NE Atlantic	250	2.46	Lusher et al., 2014
Portugal, coast	180–335	0.002–0.036	Frias et al., 2014
South. California coast	333	5–7.25	Moore et al., 2002
NW Mediterranean	333	0.116	Collignon et al., 2012
Mediterranean coast of Israel	333	7.68	van der Hal et al., 2017
Central-W Mediterr.	500	0.15	de Lucia et al., 2014
Arctic polar waters	333	0.34 \pm 0.31	Lusher et al., 2015
Southeast Bering Sea	505	<0.1	Doyle et al. (2011)
Bohai Sea, China	330	0.33 \pm 0.36	Zhang et al. (2017)
NW Pacific, Russia	100	0.13 – 1.7	This study

Table 18. Global comparison of microplastic pollution in the coastal marine water areas

In view that it is difficult to find enough data on microplastics in the tidal water (except for intertidal sediments) to make a global comparison, we compared our data for offshore samples with comparable global data collected by neuston/manta net trawling in coastal waters worldwide (table 18). The data in the Peter the Great Gulf show concentrations in the lower range of those reported in other coastal area similar to concentration levels found in the Arctic. The results are not surprising because of relatively low industrial and urban activities in the study area compared to other coastal areas in Europe and Asia.

5.7. Suggested seasonal factor and land-based sources

One of the main goals of this study was to identify possible seasonal variations and impact of land-based pollution sources on the composition and concentrations of microplastics. For several sites it was impossible to collect seasonal samples. For example, on sites 3, 4, 6, 11, and 12 only one sample per each site was collected (summer samples for sites 3, 4, and 6 and autumn samples for sites 11 and 12). For five sampling sites, 2 samples were collected in summer and in autumn (2, 5, 7, 9, and 10). At site 1, we collected 3 samples (2 in autumn and 1 in summer). At site 8, nine samples were collected (July 2016; February, May, June, July, August, September, October, and November 2017). Based on this information we analyzed seasonal differences in the microplastic pollution (where two or more seasonal samples were available) according to possible land based sources. In case of the only seasonal sample available, only land-based source impacts are discussed.

At site 1, which is exposed to a strong impact from the Tumen River, we collected three samples (sample 1 in September 2016, sample 2 in July 2017, and sample 3 in October 2017). Overall number of microplastic particles is almost similar in sample 1 and sample 2 (54 and 56 p./m³, respectively) but decreased in sample 3 (31 p./m³). The number of fibers and films is higher in summertime (fig.20, left). The amount of EPS foam is also higher during a warmer season (1 and 2). Strong increase of fragments is observed in July 2017. A possible explanation of these fluctuations is the intensity of Tumen River discharge, which decreases in the

autumn. Several additional data obtained in the lower Tumen River suggest that the increased or decreased concentrations of microplastics in the river water correspond to the increased or decreased concentrations of microplastics in the samples from marine coastal water near the estuary collected at the same time.

At site 2 no seasonal changes were observed and concentrations of microplastics were very low comparable to site 1, despite their relative proximity (table 3).

Site 1. Khasan shore

Site 3. Minonosok Inlet




Fig. 20.

left - seasonal variation of microplastic structure in the tidal water of site 1 (Khasan seashore) in particles per m^3 . (1- September 2016, 2 – July 2017, and 3 – October 2017);

right - structure of microplastic in the tidal zone of Minonosok Inlet (June 2016)

At site 3 there is an evidence that fragmentation of beached derelict aquacultural gear is a source of numerous fragments in the structure of microplastics (fig.20, right). The large amount of beached EPS also evidences that there is an increased transport of plastic litter from the Tumen River (Kozlovskii et al., 2016)

Site 4 is a remote natural protected area and unlike site 3, it is not exposed to any aquacultural activities. Therefore it is difficult to suggest possible land-based sources of pollution there (table 5).

Site 5 (a beach near the Slavyanka settlement) is a typical recreation area. The samples were collected there in late September 2016 and in late July 2017 (the time of increased recreant inflow). The summer sample was characterized by

comparatively higher concentrations of microplastics, while in late September the pollution was minimal (fig.21, upper left).

Sites 9 and 10 are also characterized by seasonal recreation impact and in this relation can be compared to site 5. Another important factor is their close vicinity to the urban area. In all cases, increased concentrations of microplastic particles are observed in summer (July - August). Another important aspect is that basic component of microplastics in the tidal zone in sites 5, 9, and 10 are fibers (fig. 21).

At site 6, only one sample was collected and total microplastic concentrations were comparatively low (1.5 p./m³). Because of that, it is difficult to suggest any impact of land sources of pollution especially considering that the territory adjoining to this site is low populated and is not used for recreation.

At site 7 we also collected two samples: in late September 2016 and in late July 2017. The sources of pollution there are difficult to identify because it is located not far from the Razdolnaya/Suifen River mouth and just opposite to the urban area of Vladivostok. Also, the site experiences some impact of small-scale fisheries (fig. 21, upper right)

The most regular monitoring was carried out in the tidal water of **Site 8** (Chaika Beach). Nine samples were collected in 2016 and 2017, including winter, spring, summer, and autumn samples (fig. 22). Varying concentration and structure of microplastic contamination are evident for this site. The lowest concentration is peculiar for winter season. Because this area is ice covered in winter, the reason for such small concentration compared to other seasons may be the ability of the ice cover to contain plastic particles. Consequently, microplastic contamination of the unfrozen water under the ice is minimal.

In spring, concentrations of microplastics considerably increase and is comparable to summer concentrations.

In autumn, the concentrations gradually decrease.

Structure of microplastics in this area evidence high contamination with fibers, both in the tidal zone and in the bay (fig. 22 and 23). It is similar to sites 5, 9, and 10. This may be a result of wastewater discharge, so there is a need for the related




Fig. 21

Upper left - seasonal variation of microplastic structure in the tidal water of site 5 (Beach near Slavyanka Settlement) in particles per m³. (1- September 2016, 2 – July 2017);

Upper right - seasonal variation of microplastic structure in the tidal water of site 7 (Peschany peninsula) in particles per m³. (1- September 2016, 2 – July 2017)

Lower left – June and August structure of microplastics in the tidal water of site 9;

Lower right - seasonal variation of microplastic structure in the tidal water of site 10

monitoring.

A study (Browne et al., 2011) evidence that machine-washing of garments may produce several thousand fibers per wash. One of suggestions indicated in this study is that late in the year the increment of in fiber concentrations from land-based sources should be evidenced up to 700 % due to washing of heavier winter clothes. In case with Chaika area, no evidence of higher concentrations of fibers was evidenced in late autumn (fig. 22).

Sites 11 and 12 were sampled only once in autumn 2017 so more observation is needed to conclude on the possible land-based impacts and seasonal distribution of microplastics in the water.




Fig. 22. Seasonal variation of microplastic structure in the tidal water of site 8 (Chaika Beach) in particles per m³.




Fig. 23. Seasonal variation of microplastic structure in the bay (neuston trawl) near site 8 (Chaika Beach) in particles per m³.

6. Conclusion

This is one of the first surveys of the microplastic coastal water pollution in the Russian part of NOWPAP area, and there are several findings.

Some general conclusion can be drawn that the microplastic pollution issue is actual for the southern Far East of Russia, though its scale is evidently lower than in the neighboring NOWPAP countries.

During this survey we identified the most important local hotspots of microplastic pollution, including the Tumen River estuary area as the most outstanding, and to a lesser extent coastal water in the vicinity of Vladivostok agglomeration (inner part of the Amur Bay water area).

Based on the seasonal changes and structure of microplastic particles from different sampling sites we can identify the most evident land-based sources of the contamination. For example, river discharge in case with the Tumen River estuary area, or coastal degradation of derelict domestic waste or fishing/aquacultural gear on remote shores which lack regular cleanups, for example in the Minonosok Inlet. Within urban and recreational areas there is an evidence of higher concentrations of fibers. In case with recreational locations, which are exposed to an increased anthropogenic load in summer, total summer concentrations of microplastics and

especially fibers are higher. In case with urban areas, where fibers are the basic morphological type of microplastics, possible source of the contamination is domestic wastewater discharge (after machine washing). Due to this fact, additional survey of the effluent is necessary.

In general, it is evident that further survey in the coastal marine area of southern Far East of Russia is necessary with a detailed assessment of suggested land-based pollution sources. Particularly, specific attention should be paid to the river discharge and urban wastewater.

It is also necessary to assess possible negative impacts on marine and freshwater biota in the Russian part of NOWPAP.

7. Acknowledgement

We thank NOWPAP member countries for financial support and express our cordial gratitude to our distinguished colleagues from the NOWPAP member countries who assisted us in developing methods for this study and teaching us their national procedures of microplastic monitoring.

In particular, we would like to thank Dr. Sang Hee Hong and her colleagues from Korean Institute of Ocean Science and Technology (KIOST) who provided the protocols for water sample treatment and FTIR analysis.

We thank Dr. Keiichi Uchida and his colleagues from Tokyo University of Marine Science and Technology (TUMSAT) who invited us to participate in a joint marine litter research cruise in the Japanese coastal waters and provided field training for microplastic sampling.

And, of course, special thanks are to Dr. Atsuhiko Isobe and his colleagues from Kyushu University who provided us the microplastic assessment methodology used in Japan.

References

U. Aytan, A. Valente, Y. Senturk, R. Usta, F. Basak, E. Sahin, R. E. Mazlum, E. Agirbas // First evaluation of neustonic microplastics in Black Sea waters // Marine Environmental Research, Volume 119, August 2016, pp. 22-30

Y. Y. Blinovskaya // Microplastic is the macroproblem of the World Ocean including Russian Far East //

http://www.npec.or.jp/NWPacific_node/report/document/Microplastics%20is%20the%20macroproblem%20of%20the%20world%20ocean%20including%20russian%20far%20east.pdf

M.A. Browne, P. Crump, S. J. Niven, E. Teuten, A. Tonkin, T. Galloway, R. Thompson //

Accumulation of microplastic on shorelines worldwide: sources and sinks. (2011).

Environ. Sci. Technol., 45: 9175 – 9179

P. Davison, R.G. Asch (2011) Plastic ingestion by mesopelagic fishes in the North Pacific Subtropical Gyre. Mar Ecol Prog Ser 432:173-180

J.P.W. Desforges, M. Galbraith, N. Dangerfield, P.S. Ross // Widespread distribution of microplastics in subsurface seawater in the NE Pacific Ocean // Mar. Pollut. Bull., 79 (2014), pp. 94-99

M. J. Doyle, W. Watson, N. M. Bowlin, S. B. Sheavly // Plastic particles in coastal pelagic ecosystems of the Northeast Pacific ocean // Marine Environmental Research 71 (2011) pp. 41-52

J.P.G.L. Frias, V. Otero, P. Sobral // Evidence of microplastics in samples of zooplankton from Portuguese coastal waters // Mar. Environ. Res., 95 (2014), pp. 89-95

N. van der Hal, A. Ariel, D. L. Angel // Exceptionally high abundances of microplastics in the oligotrophic Israeli Mediterranean coastal waters // Marine Pollution Bulletin 116 (2017), pp. 151–155

V. Hidalgo-Ruz, L. Gutow, R. C. Thompson, M. Thiel // Microplastics in the Marine Environment: A Review of the Methods Used for Identification and Quantification // *Environ. Sci. Technol.*, 2012, 46 (6), pp 3060–3075

J.-H. Kang, O. Y. Kwon, K.-W. Lee, Y. K. Song, W. J. Shim // Marine neustonic microplastics around the southeastern coast of Korea // Marine Pollution Bulletin, Volume 96, Issues 1–2, 15 July 2015, pp. 304-312

N.V. Kozlovskii, S.H. Hong, Y.K. Song, A.N. Kachur // Distribution of beached marine litter and floating microplastics in the Minonosok Inlet of Possiet Bay of the Peter the Great Gulf // 2016

http://www.npec.or.jp/NWPacific_node/report/document/Distribution%20of%20beached%20marine%20litter%20and%20floating%20microplastics%20in%20the%20minonosok%20inlet%20of%20possiet%20bay%20of%20the%20peter%20the%20great%20gulf.pdf

Laboratory Methods for the Analysis of Microplastics in the Marine Environment: Recommendations for quantifying synthetic particles in waters and sediments NOAA Marine Debris Program National Oceanic and Atmospheric Administration U.S. Department of Commerce Technical Memorandum NOS-OR&R-48 July 2015 //

https://marinedebris.noaa.gov/sites/default/files/publications-files/noaa_microplastics_methods_manual.pdf

A.L. Lusher, A. Burke, I. O'Connor, R. Officer // Microplastic pollution in the northeast Atlantic Ocean: validated and opportunistic sampling // Mar. Pollut. Bull., 88 (2014), pp. 325-333,

A.L. Lusher, V. Tirelli, I. O'Connor, R. Officer // Microplastics in Arctic polar waters: the first reported values of particles in surface and sub-surface samples. // Scientific reports 5, article # 14947

<https://www.nature.com/articles/srep14947>

C.J. Moore, S.L. Moore, S.B. Weisberg, G.L. Lattin, A.F. Zellers // A comparison of neustonic plastic and zooplankton abundance in southern California's coastal waters//

Mar. Pollut. Bull., 44 (2002), pp. 1035-1038

F. Murray, P.R. Cowie // Plastic contamination in the decapod crustacean *Nephrops norvegicus* (Linnaeus, 1758). Mar. Pollut. Bull. (2011), doi:10.1016/j.marpolbul.2011.03.032

F. Norén // Small Plastic Particles in Coastal Swedish Waters. N-Research Report, Commissioned by KIMO Sweden (2008) 11 pp

M. D. Robards, J. F. Piatt, and K. D. Wohl. 1995. Increasing frequency of plastic particle ingestion by seabirds in the subarctic North Pacific. Marine Pollution Bulletin 30: 151-157.

R. Sussarellu, M. Suquet, Y. Thomas, Ch. Lambert, C. Fabioux, M. E. J. Pernet, N. L. Goïc, V. Quillien, Ch. Mingant, Y. Epelboin, Ch. Corporeau, J. Guyomarch, J. Robbins, I. Paul-Pont, P. Soudant, and A. Huvet (2016). Oyster reproduction is affected by exposure to polystyrene microplastics. Proc Natl Acad Sci U S A.

113(9): 2430–2435.

E. L. Teuten, J. M. Saquing, D. R. Knappe, U. Barlaz, M. A. Jonsson, S. Björn, A. Rowland, S. J. Thompson, R. C. Galloway, T. S. Yamashita, R. Ochi, D. Watanuki, Y. Moore, C. Viet, P. Tana, T. S. Prudente, M. Boonyatumanond, R. Zakaria, M. P. Akkhavong, K. Ogata, Y. Hirai, H. Iwasa, S. Mizukawa, K. Hagino, Y. Imamura, A. Saha, M. Takada, S. (2009) Transport and release of chemicals from plastics to the environment and to wildlife. *Philosophical Transactions of the Royal Society B* 364, 2027-2045.

A. L. Yakimenko, Y. Y. Blinovskaya // results of microplastic monitoring in the coastal marine area of the southern Primorsky Krai // Scientific magazine 'Concept' – 2016. – V. 11. – pp. 3576–3580. – URL: <http://e-koncept.ru/2016/86753.htm>. (in Russian)

W. Zhang, S. Zhang, J. Wang, Y. Wang, J. Mu, P. Wang, X. Lin, D. Ma // Microplastic pollution in the surface waters of the Bohai Sea, China // *Environmental Pollution*, Volume 231, Part 1, December 2017, pp. 541-548

S. Zhao, L. Zhu, T. Wang, D. Li // Suspended microplastics in the surface water of the Yangtze Estuary System, China: first observations on occurrence, distribution // *Mar. Pollut. Bull.*, 86 (2014), pp. 562-568

Технический отчет

**Microplastic pollution in the coastal water
of the Peter the Great Gulf: content and distribution.
The first stage of survey**

**Загрязнение прибрежных вод
залива Петра Великого микропластиком: его состав и распределение.
Первый этап исследований**

На английском языке

Авторы: Н. В. Козловский, А.Н. Качур.
Отв. Редактор: С.Ю. Монинец

Отпечатано с оригинал-макета,
подготовленного в Тихоокеанском институте географии ДВО РАН

Формат 60x84/8. Усл. п.л 5,12. Уч.-изд. л. 4,9.
Тираж 500 экз. Заказ 2

ФГБУН Тихоокеанский Институт Географии
Дальневосточного отделения Российской Академии Наук
690041, Владивосток, ул. Радио, 7

Отпечатано в Информационно-полиграфическом хозрасчетном
центре ТИГ ДВО РАН
690041, Владивосток, ул. Радио, 7